

ANDROSTENEDIONE,  
SERUM

## Orderable - ANDRO

Turnaround Time: 10 days

### Specimen:

Adult	Pediatric
6 mL Red top Vacutainer tube	0-2 years: 2 x 0.5 mL Red top Microtainers 2-10 years: 3 mL Red top Vacutainer tube
<b>Avoid blood collection tubes with separator gels (Gold top Vacutainer tubes)</b>	
Light Green (Li-Heparin) or Lavender (EDTA) top tubes are <b>NOT</b> acceptable	



**Laboratory:**  
Toxicology/Special  
Chemistry



**Requisition:**  
GENERAL LABORATORY  
REQUISITION



**Method of Analysis:**  
LC-MS/MS

### Collection Information:

Minimum volume of serum required is 0.5 mL.

### Reference Ranges:

Age	Reference Interval (nmol/L)
0 - <14 days	≤2.5
14 days - <1 year	≤2.1
1 - <6 years	<1.0
6 - <10 years	<1.0
10 - <12 years	≤2.5



**Test Schedule:**  
Monday-Friday, 0800-  
1600

Male	
Age	Range
12 - <15 years	≤2.0
15 - <19 years	≤3.6
19 - <31 years	1.8-7.7
31 - <51 years	1.4-6.6
51 - <89 years	1.8-7.7

ANDROSTENEDIONE,  
SERUM

Female	
Age	Range
12 - <15 years	≤6.0
15 - <19 years	≤6.5
Follicular Phase:	1.2-8.7
Midcycle:	2.1-10.0
Luteal Phase:	1.1-8.2
Post-menopausal:	≤2.6

**Storage and Shipment:**

Store and ship serum frozen.