

IMMUNOGLOBULIN  
A, PLASMA/  
SERUM

## Orderable - IGAM

Turnaround Time: 4 hours

### Alternate Name(s):

IgA

### Specimen:

Adult	Pediatric
4.5 mL Light Green top Vacutainer tube	0-2 years: Red 0.5 mL Microtainer 2-10 years: 2 mL Red top tube
5 mL Gold top Vacutainer tube or 6 mL Red top Vacutainer tube are also acceptable	

### Collection Information:

Fasting specimen preferred.

Avoid hemolysis.

Turbidimetric assays are not suitable for measurement of highly lipemic or hemolytic samples or samples containing high levels of circulating immune complexes

### Reference Ranges:

Age	Range
<1 year:	0.0-0.1 g/L
1-<3 years:	0.0-0.8 g/L
3-<6 years:	0.1-1.4 g/L
6-<14 years:	0.3-2.2 g/L
14-<19years:	0.4-2.9 g/L
Adult:	0.7-4.0 g/L

### Comments:

Serum IgG, IgA, and IgM are tested simultaneously as a group.



**Laboratory:**  
Core Lab



**Requisition:**  
GENERAL LABORATORY  
REQUISITION



**Method of Analysis:**  
Immunoturbidimetric  
Assay



**Test Schedule:**  
Daily