

MYOSITIS ANTIBODIES PROFILE

Orderable - .MYOAP

Turnaround Time: 5 days

Alternate Name(s):

Autoimmune Myositis Panel
Myositis-specific antibodies
Anti-Mi-2 alpha
Anti-Mi-2 beta
Anti-TIF1 gamma
Anti-MDA5
Anti-NXP2
Anti-SAE1
Anti-Ku
Anti-PM-Scl 100
Anti-PM-Scl 75
Anti-JO1
Anti-PL-7
Anti-PL-12
Anti-EJ
Anti-OJ
Anti-Ro-52
Anti-Ha
Anti-Ks
Anti-Zo Anti-cN1A
Anti-SRP



Laboratory:
Clinical Immunology Lab



Requisition:
[CLINICAL IMMUNOLOGY
REQUISITION](#)



Method of Analysis:
Immunoblot



Test Schedule:
Batch analysis

Specimen:


Adult	Pediatric
5 mL Gold top Vacutainer tube or 6 mL Red top Vacutainer tube	0-2 years: Red 0.5 Microtainer 2-10 years: 2 mL Red top
EDTA, Heparin or Citrate plasma are also acceptable	

Collection Information:

Minimum volume of serum or plasma required for testing is 0.5 mL.

Reference Ranges:

Negative



MYOSITIS ANTIBODIES PROFILE

Interpretive Comments:

Immunoblot results will be reported as either negative or positive for specific antibodies as the following.

- 1+ Weak Positive
- 2+ Positive
- 3+ Strong Positive

Comments:

This test is available exclusively to Rheumatologists and Respiriologists at LHSC/SJH and accepted from referred in locations.

Sample Stability:

Refrigerated: 14 days
Frozen: 30 days