



Pathology and Laboratory Medicine

SELENIUM, PLASMA Orderable - SEP

Turnaround Time: 10 days

Specimen:

6 mL K2-EDTA Royal Blue Vacutainer tube

Collection Information:

Trace Elements Collection Requirements



Requisition: TRACE ELEMENTS

Trace Elements Lab

Laboratory:

REQUISITION



Method of Analysis: ICP-MS/MS



Test Schedule:Batched analysis

Reference Ranges:

SI Units (Reported on Patient Chart):	
Age	Reference Range
0-12 months:	0.72 -1.21 μmol/L
1-5 years:	1.22-1.82 μmol/L
6- 9 years:	1.28-2.04 μmol/L
≥10 years:	1.33-2.03 μmol/L

Conventional Units:	
Age	Reference Range
0-12 months:	56.9-95.6 μg/L
1- 5 years:	96.4-143.8 μg/L
6-9 years:	100.8-161.4 μg/L
≥10 years:	105.3-160.4 μg/L

Concentration of Selenium is much higher in erythrocytes than in plasma or serum. The results of these elements in plasma or serum may be falsely elevated if not separated within 30 minutes and/or hemolysis is present.

Reference Ranges are based on Non-Occupationally exposed population.

Interpretive Comments:

Find Interpretive Comment and Clinical Information here:







Pathology and Laboratory Medicine

Specimen Stability:

Ambient: 22 days

Refrigerated: 14 months Frozen: 14 months