

APPENDIX D

PRECAUTIONS *DURING AGMPs* in CRITICAL CARE

Precautions A	Precautions B	Precautions C
PROTECTED Droplet + Contact + Enhanced <ul style="list-style-type: none"> Fit-tested N95 Respirator Disposable goggles Full face shield Hair Bouffant Level 3 gown or higher Extended cuff nitrile gloves or sterile gloves that cover wrists 	Droplet + Contact + Enhanced <ul style="list-style-type: none"> Fit-tested N95 Respirator Full face shield Level 2 gown or higher Gloves 	Droplet + Contact <ul style="list-style-type: none"> Procedure/surgical mask Protective eyewear (e.g. attached visor or face shield) Long-sleeved gown (Level 2 if low to moderate fluid exposure risk) Gloves
Negative pressure (if available)		

Following Aerosol Generating Procedures:

PPE must be worn for 30 minutes after completion of an AGMP. AGMP PPE should be worn in rooms where there is a likelihood of AGMPs being performed.

Aerosol Generating Medical Procedures (AGMP)	Precaution Category: Failed ARI AND COVID-19 pending * testing not indicated on outpatients OR COVID-19 positive OR Unable to assess OR Household contact of confirmed case	Precaution Category: COVID-19 negative NOT suspect COVID-19
Intubation	A	A
Bronchoscopy (including percutaneous tracheostomy)	A	A
CPR	A	A
Extubation	B (self-assess for +goggles +bouffant)	C
Bag-valve-mask ventilation	B (with filter, self-assess for +goggles+bouffant)	C
Open Circuit/Airway Suctioning	B	C
CPAP/ NIV/High Flow Nasal Oxygen	B (negative pressure room if available)	C
Ventilator or circuit change	B (clamp ETT momentarily)	C
Tracheostomy tube change/decannulation	B (self-assess for + goggles + bouffant)	C
Nebulizer treatments	B (meter-dosed with spacer recommended)	C
Chest tube insertion with potential air leak	B (Level 3 gown and sterile gloves as part of sterile procedure; + goggles +bouffant)	C
Sputum induction, cough-assist, breath stacking	B (+ goggles + bouffant)	C
All upper GI procedures	B	C
Other endoscopic procedures	B	C Routine Practices