

IMIPRAMINE/
DESIPRAMINE,
SERUM/
PLASMA

Orderable - IMIP

Turnaround Time: 7 days

Alternate Name(s):

Tofranil

Specimen:

6 mL Red top Vacutainer tube

Lavender or Pink (EDTA) top tubes are also acceptable

**Avoid blood collection tubes with separator gels (e.g., Gold top).
Light Green (Li-Heparin) top tubes are also NOT acceptable.**

Collection Information:

Minimum volume of serum or EDTA plasma required is 3 mL.

To monitor therapy, draw trough specimen prior to morning dose or 10 - 12 h after last drug administration. Separate and freeze as soon as possible.

Reference Ranges:

Please note change in units, therapeutic range, and toxic range as of October 18, 2022.

Imipramine + Desipramine:	175-300 ng/mL
------------------------------	---------------

Critical Value:

Toxic:

Imipramine + Desipramine:	> 300 ng/mL
------------------------------	-------------



Laboratory:
Core Lab



Requisition:
GENERAL LABORATORY
REQUISITION



Method of Analysis:
LC-MS/MS



Test Schedule:
Referred out Monday-
Thursday



Referred Out Location:
[In-Common Laboratories](#)

IMIPRAMINE/
DESIPRAMINE,
SERUM/
PLASMA

Storage and Shipment:

Send frozen sample ASAP